

# Migratory connectivity analysis

by EURING Migration Atlas


*Uria aalge* (EURING code 06340)

## 1.1 Connectivity between individuals

The analysis evaluated 1490 individuals (2980 encounters) filtered from a total of 97725 records in the EURING databank which were considered for the Atlas. The species shows a significant connectivity from pattern transference (Table 06340-1; Figure 06340-1).

**Table 06340-1.** Results from the migratory connectivity analysis. For each cluster, the degree of connectivity ( $r_M$ ), its statistical significance (p-value) and 95% confidence interval limits are shown. When the p-value is less than or equal to 0.1, the degree of clustering structure (oasw) and the best number of clusters identified are reported.

Cluster name	Level of clustering	N individuals	Migratory connectivity ( $r_M$ )	p-value	Lower 95% confidence limit	Upper 95% confidence limit	Best number of clusters	oasw
0	0	1490	0.366	0.001	0.329	0.407	2	0.431




**Figure 06340-1.** Map showing 95% kernel contours of of first-level clusters identified by the migratory connectivity analysis, if any, or 95% kernel contours of all encounters, in case of no clustering structure. Solid lines indicate the clusters in the breeding range, dotted lines those in the non-breeding range. Different contour colours correspond to different clusters, as reported in legend. The species distribution range is also shown (breeding range: blue; non-breeding range: dark grey; resident range: beige; from BirdLife International, 2019).


## 1.2 Sensitivity analysis

Results of power analysis and validation. Analyses at the species level were re-run on subsamples of individuals of decreasing size (100 repetitions per subsample size), according to simple random sampling of individuals (Figure 06340-2) and stratified sampling of individuals within the breeding range (Figure 06340-3) and the non breeding range (Figure 06340-4). For stratified sampling, we selected individuals with a

probability inversely proportional to the number of observation in each country. Figures below report the results of the procedure.




**Figure 06340-2.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis (i.e. proportion of times the analyses on the subset of individuals was significant). Bottom left: Proportion of times the analysis provides the observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.




**Figure 06340-3.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis. Bottom left: Proportion of times the analysis provides the

observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.



**Figure 06340-4.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis. Bottom left: Proportion of times the analysis provides the observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.

The comparison between the bootstrapped distribution of  $r_M$  values from live recaptures and dead recoveries is significant ( $p = 0.013$ ); Figure 06340-5).



**Figure 06340-5.** Comparison between the bootstrapped distributions of connectivity value for alive recaptures (filling lines with angle=45°) and dead recoveries (filling lines with angle=375°).


## 2. Connectivity between pre-defined regions

The species shows low connectivity ( $MC = 0.033$ ;  $MC = 0.032$  when adjusted for absolute abundance) between 4 breeding regions and 6 non breeding regions (Table 06340-2; Figure 06340-6).

**Table 06340-2.** Transition probabilities between pre-defined regions. Estimated abundance (number of individuals) in each breeding region is also reported.

Breeding region	Abundance	Non breeding region	Transition probability
Central Europe	4448	Central Europe	0.373
Central Europe	4448	North Europe	0.382
Central Europe	4448	North-west Europe	0.078
Central Europe	4448	West Europe	0.167
North Europe	415670	East Europe	0.022
North Europe	415670	North Europe	0.946
North Europe	415670	North-west Europe	0.022
North Europe	415670	West Europe	0.011
North-west Europe	2274804	Central Europe	0.031
North-west Europe	2274804	North Europe	0.369
North-west Europe	2274804	North-west Europe	0.297
North-west Europe	2274804	South-west Europe	0.038
North-west Europe	2274804	West Europe	0.265

Breeding region	Abundance	Non breeding region	Transition probability
West Europe	586	North-west Europe	0.333
West Europe	586	West Europe	0.667



**Figure 06340-6.** Map showing pre-defined regions in different colours, with black arrows linking centroids of individual encounters in different regions. Arrow width is proportional to transition probability.

## Reference

BirdLife International and Handbook of the Birds of the World (2019). Bird species distribution maps of the world. Version 2019.1. Available at <http://datazone.birdlife.org/species/requestdis>.