

Migratory connectivity analysis

by EURING Migration Atlas

Alca torda (EURING code 06360)

1.1 Connectivity between individuals

The analysis evaluated 521 individuals (1042 encounters) filtered from a total of 28889 records in the EURING databank which were considered for the Atlas. The species shows a significant connectivity from clustering, with a number of first-level clusters = 2 (Table 06360-1; Figure 06360-1).

Table 06360-1. Results from the migratory connectivity analysis. For each cluster, the degree of connectivity (r_M), its statistical significance (p-value) and 95% confidence interval limits are shown. When the p-value is less than or equal to 0.1, the degree of clustering structure (oasw) and the best number of clusters identified are reported.

Cluster name	Level of clustering	N individuals	Migratory connectivity (r_M)	p-value	Lower 95% confidence limit	Upper 95% confidence limit	Best number of clusters	oasw
0	0	521	0.485	0.001	0.436	0.536	2	0.565
1	1	400	0.183	0.001	0.122	0.246	2	0.506
2	1	121	0.341	0.001	0.231	0.475	3	0.581
11	2	249	0.274	0.001	0.154	0.387	7	0.399
12	2	151	0.042	0.186	-0.052	0.183	-	-
21	2	6	-	-	-	-	-	-
22	2	13	-	-	-	-	-	-
23	2	102	0.011	0.350	-0.047	0.129	-	-




Figure 06360-1. Map showing 95% kernel contours of of first-level clusters identified by the migratory connectivity analysis, if any, or 95% kernel contours of all encounters, in case of no clustering structure. Solid lines indicate the clusters in the breeding range, dotted lines those in the non-breeding range. Different contour colours correspond to different clusters, as reported in legend. The species distribution range is also shown (breeding range: blue; non-breeding range: dark grey; resident range: beige; from BirdLife International, 2019).

1.2 Sensitivity analysis

Results of power analysis and validation. Analyses at the species level were re-run on subsamples of individuals of decreasing size (100 repetitions per subsample size), according to simple random sampling of individuals (Figure 06360-2) and stratified sampling of individuals within the breeding range (Figure 06360-3) and the non breeding range (Figure 06360-4). For stratified sampling, we selected individuals with a

probability inversely proportional to the number of observation in each country. Figures below report the results of the procedure.




Figure 06360-2. Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis (i.e. proportion of times the analyses on the subset of individuals was significant). Bottom left: Proportion of times the analysis provides the observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.




Figure 06360-3. Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis. Bottom left: Proportion of times the analysis provides the

observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.




Figure 06360-4. Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis. Bottom left: Proportion of times the analysis provides the observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.

The comparison between the bootstrapped distribution of r_M values from live recaptures and dead recoveries is not significant ($p = 0.08$); Figure 06360-5).




Figure 06360-5. Comparison between the bootstrapped distributions of connectivity value for alive recaptures (filling lines with angle=45°) and dead recoveries (filling lines with angle=375°).

2. Connectivity between pre-defined regions

The species shows moderate connectivity (MC = 0.408; MC = 0.407 when adjusted for absolute abundance) between 4 breeding regions and 7 non breeding regions (Table 06360-2; Figure 06360-6).

Table 06360-2. Transition probabilities between pre-defined regions. Estimated abundance (number of individuals) in each breeding region is also reported.

Breeding region	Abundance	Non breeding region	Transition probability
East Europe	12511	Central Europe	0.667
East Europe	12511	East Europe	0.333
North Europe	138400	Central Europe	0.342
North Europe	138400	East Europe	0.202
North Europe	138400	North Europe	0.421
North Europe	138400	North-west Europe	0.009
North Europe	138400	West Europe	0.026
North-west Europe	837525	Central Europe	0.025
North-west Europe	837525	North Africa	0.037
North-west Europe	837525	North Europe	0.062
North-west Europe	837525	North-west Europe	0.347
North-west Europe	837525	South-west Europe	0.342
North-west Europe	837525	West Europe	0.186

Breeding region	Abundance	Non breeding region	Transition probability
West Europe	76	West Europe	1.000




Figure 06360-6. Map showing pre-defined regions in different colours, with black arrows linking centroids of individual encounters in different regions. Arrow width is proportional to transition probability.

Reference

BirdLife International and Handbook of the Birds of the World (2019). Bird species distribution maps of the world. Version 2019.1. Available at <http://datazone.birdlife.org/species/requestdis>.