

# Migratory connectivity analysis

by EURING Migration Atlas


*Prunella modularis* (EURING code 10840)

## 1.1 Connectivity between individuals

The analysis evaluated 10936 individuals (21872 encounters) filtered from a total of 507605 records in the EURING databank which were considered for the Atlas. The species shows a significant connectivity from clustering, with a number of first-level clusters = 2 (Table 10840-1; Figure 10840-1).

**Table 10840-1.** Results from the migratory connectivity analysis. For each cluster, the degree of connectivity ( $r_M$ ), its statistical significance (p-value) and 95% confidence interval limits are shown. When the p-value is less than or equal to 0.1, the degree of clustering structure (oasw) and the best number of clusters identified are reported.

Cluster name	Level of clustering	N individuals	Migratory connectivity ( $r_M$ )	p-value	Lower 95% confidence limit	Upper 95% confidence limit	Best number of clusters	oasw
0	0	10936	0.960	0.001	0.949	0.971	2	0.519
1	1	1689	0.706	0.001	0.650	0.765	3	0.491
2	1	9247	0.991	0.001	0.983	0.997	9	0.457




**Figure 10840-1.** Map showing 95% kernel contours of of first-level clusters identified by the migratory connectivity analysis, if any, or 95% kernel contours of all encounters, in case of no clustering structure. Solid lines indicate the clusters in the breeding range, dotted lines those in the non-breeding range. Different contour colours correspond to different clusters, as reported in legend. The species distribution range is also shown (breeding range: blue; non-breeding range: dark grey; resident range: beige; from BirdLife International, 2019).


## 1.2 Sensitivity analysis

Results of power analysis and validation. Analyses at the species level were re-run on subsamples of individuals of decreasing size (100 repetitions per subsample size), according to simple random sampling of individuals (Figure 10840-2) and stratified sampling of individuals within the breeding range (Figure 10840-3) and the non breeding range (Figure 10840-4). For stratified sampling, we selected individuals with a

probability inversely proportional to the number of observation in each country. Figures below report the results of the procedure.




**Figure 10840-2.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis (i.e. proportion of times the analyses on the subset of individuals was significant). Bottom left: Proportion of times the analysis provides the observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.




**Figure 10840-3.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis. Bottom left: Proportion of times the analysis provides the

observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.



**Figure 10840-4.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis. Bottom left: Proportion of times the analysis provides the observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.

The comparison between the bootstrapped distribution of  $r_M$  values from live recaptures and dead recoveries is significant ( $p < 0.001$ ); Figure 10840-5).



**Figure 10840-5.** Comparison between the bootstrapped distributions of connectivity value for alive recaptures (filling lines with angle=45°) and dead recoveries (filling lines with angle=375°).


## 2. Connectivity between pre-defined regions

The species shows moderate/high connectivity (MC = 0.642; MC = 0.642 when adjusted for absolute abundance) between 5 breeding regions and 7 non breeding regions (Table 10840-2; Figure 10840-6).

**Table 10840-2.** Transition probabilities between pre-defined regions. Estimated abundance (number of individuals) in each breeding region is also reported.

Breeding region	Abundance	Non breeding region	Transition probability
Central Europe	5702700	Central Europe	0.759
Central Europe	5702700	South-central Europe	0.043
Central Europe	5702700	South-west Europe	0.149
Central Europe	5702700	West Europe	0.050
North Europe	3001000	Central Europe	0.094
North Europe	3001000	North Europe	0.189
North Europe	3001000	South-central Europe	0.057
North Europe	3001000	South-east Europe	0.019
North Europe	3001000	South-west Europe	0.415
North Europe	3001000	West Europe	0.226
North-west Europe	9840630	North-west Europe	1.000
South-west Europe	2644423	South-west Europe	1.000
West Europe	3428521	South-west Europe	0.003

Breeding region	Abundance	Non breeding region	Transition probability
West Europe	3428521	West Europe	0.997



**Figure 10840-6.** Map showing pre-defined regions in different colours, with black arrows linking centroids of individual encounters in different regions. Arrow width is proportional to transition probability.

## Reference

BirdLife International and Handbook of the Birds of the World (2019). Bird species distribution maps of the world. Version 2019.1. Available at <http://datazone.birdlife.org/species/requestdis>.