

# Migratory connectivity analysis

by EURING Migration Atlas

*Serinus serinus* (EURING code 16400)


## 1.1 Connectivity between individuals

The analysis evaluated 672 individuals (1344 encounters) filtered from a total of 33377 records in the EURING databank which were considered for the Atlas. The species shows a significant connectivity from clustering, with a number of first-level clusters = 2 (Table 16400-1; Figure 16400-1).

**Table 16400-1.** Results from the migratory connectivity analysis. For each cluster, the degree of connectivity ( $r_M$ ), its statistical significance (p-value) and 95% confidence interval limits are shown. When the p-value is less than or equal to 0.1, the degree of clustering structure (oasw) and the best number of clusters identified are reported.

Cluster name	Level of clustering	N individuals	Migratory connectivity ( $r_M$ )	p-value	Lower 95% confidence limit	Upper 95% confidence limit	Best number of clusters	oasw
0	0	672	0.851	0.001	0.809	0.888	2	0.724
1	1	119	0.646	0.001	0.524	0.770	3	0.691
2	1	553	0.814	0.001	0.723	0.904	9	0.685
11	2	38	-0.038	0.681	-0.058	0.153	-	-
12	2	19	-	-	-	-	-	-
13	2	62	0.981	0.001	0.834	1.000	2	0.930
21	2	6	-	-	-	-	-	-
22	2	41	0.700	0.001	0.462	0.882	9	0.639
23	2	101	0.935	0.001	0.842	1.000	9	0.759
24	2	199	0.363	0.016	0.165	0.863	9	0.807
25	2	34	0.777	0.001	0.642	0.949	8	0.752
26	2	81	0.990	0.001	0.962	1.000	9	0.934
27	2	66	0.923	0.001	0.782	1.000	9	0.940
28	2	9	-	-	-	-	-	-
29	2	16	-	-	-	-	-	-
131	3	1	-	-	-	-	-	-
132	3	61	0.912	0.001	0.806	1.000	8	0.913
221	3	15	-	-	-	-	-	-
222	3	9	-	-	-	-	-	-
223	3	9	-	-	-	-	-	-
224	3	1	-	-	-	-	-	-
225	3	2	-	-	-	-	-	-
226	3	1	-	-	-	-	-	-
227	3	1	-	-	-	-	-	-
228	3	2	-	-	-	-	-	-
229	3	1	-	-	-	-	-	-
231	3	10	-	-	-	-	-	-

Cluster name	Level of clustering	N individuals	Migratory connectivity ( $r_M$ )	p-value	Lower 95% confidence limit	Upper 95% confidence limit	Best number of clusters	oasw
232	3	5	-	-	-	-	-	-
233	3	16	-	-	-	-	-	-
234	3	22	1.000	0.001	1.000	1.000	3	0.955
235	3	1	-	-	-	-	-	-
236	3	17	-	-	-	-	-	-
237	3	18	-	-	-	-	-	-
238	3	7	-	-	-	-	-	-
239	3	5	-	-	-	-	-	-
241	3	1	-	-	-	-	-	-
242	3	17	-	-	-	-	-	-
243	3	1	-	-	-	-	-	-
244	3	143	0.898	0.001	0.866	0.991	9	0.955
245	3	24	1.000	0.045	1.000	1.000	2	0.958
246	3	5	-	-	-	-	-	-
247	3	5	-	-	-	-	-	-
248	3	2	-	-	-	-	-	-
249	3	1	-	-	-	-	-	-
251	3	1	-	-	-	-	-	-
252	3	7	-	-	-	-	-	-
253	3	1	-	-	-	-	-	-
254	3	16	-	-	-	-	-	-
255	3	6	-	-	-	-	-	-
256	3	1	-	-	-	-	-	-
257	3	1	-	-	-	-	-	-
258	3	1	-	-	-	-	-	-
261	3	3	-	-	-	-	-	-
262	3	1	-	-	-	-	-	-
263	3	4	-	-	-	-	-	-
264	3	5	-	-	-	-	-	-
265	3	3	-	-	-	-	-	-
266	3	12	-	-	-	-	-	-
267	3	45	-	-	-	-	-	-
268	3	3	-	-	-	-	-	-
269	3	5	-	-	-	-	-	-
271	3	2	-	-	-	-	-	-
272	3	48	-	-	-	-	-	-
273	3	4	-	-	-	-	-	-
274	3	2	-	-	-	-	-	-
275	3	2	-	-	-	-	-	-
276	3	4	-	-	-	-	-	-
277	3	2	-	-	-	-	-	-
278	3	1	-	-	-	-	-	-
279	3	1	-	-	-	-	-	-




**Figure 16400-1.** Map showing 95% kernel contours of of first-level clusters identified by the migratory connectivity analysis, if any, or 95% kernel contours of all encounters, in case of no clustering structure. Solid lines indicate the clusters in the breeding range, dotted lines those in the non-breeding range. Different contour colours correspond to different clusters, as reported in legend. The species distribution range is also shown (breeding range: blue; non-breeding range: dark grey; resident range: beige; from BirdLife International, 2019).


## 1.2 Sensitivity analysis

Results of power analysis and validation. Analyses at the species level were re-run on subsamples of individuals of decreasing size (100 repetitions per subsample size), according to simple random sampling of individuals (Figure 16400-2) and stratified sampling of individuals within the breeding range (Figure 16400-3) and the non breeding range (Figure 16400-4). For stratified sampling, we selected individuals with a

probability inversely proportional to the number of observation in each country. Figures below report the results of the procedure.




**Figure 16400-2.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis (i.e. proportion of times the analyses on the subset of individuals was significant). Bottom left: Proportion of times the analysis provides the observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.




**Figure 16400-3.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis. Bottom left: Proportion of times the analysis provides the

observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.



**Figure 16400-4.** Top left: simulated distribution (boxplots) and observed value (red line) of connectivity. Top right: Simulated power of the analysis. Bottom left: Proportion of times the analysis provides the observed best number of cluster. Bottom right: simulated distribution (boxplots) and observed value (red line) of clustering intensity.

The comparison between the bootstrapped distribution of  $r_M$  values from live recaptures and dead recoveries is significant ( $p < 0.001$ ); Figure 16400-5).



**Figure 16400-5.** Comparison between the bootstrapped distributions of connectivity value for alive recaptures (filling lines with angle=45°) and dead recoveries (filling lines with angle=375°).

## 2. Connectivity between pre-defined regions

The species shows high connectivity (MC = 0.869; MC = 0.869 when adjusted for absolute abundance) between 5 breeding regions and 4 non breeding regions (Table 16400-2; Figure 16400-6).

**Table 16400-2.** Transition probabilities between pre-defined regions. Estimated abundance (number of individuals) in each breeding region is also reported.

Breeding region	Abundance	Non breeding region	Transition probability
Central Europe	2917200	South-central Europe	0.339
Central Europe	2917200	South-west Europe	0.643
Central Europe	2917200	West Europe	0.018
South-central Europe	3289908	South-central Europe	1.000
South-east Europe	3359560	South-east Europe	1.000
South-west Europe	42021625	South-west Europe	1.000
West Europe	432419	South-west Europe	1.000



**Figure 16400-6.** Map showing pre-defined regions in different colours, with black arrows linking centroids of individual encounters in different regions. Arrow width is proportional to transition probability.

## Reference

BirdLife International and Handbook of the Birds of the World (2019). Bird species distribution maps of the world. Version 2019.1. Available at <http://datazone.birdlife.org/species/requestdis>.